

Study of color suppressed modes $B^0 \rightarrow \bar{D}^{(*)0}\eta^{(\prime)}$

Cai-Dian Lü

CCAST (World Laboratory), P.O. Box 8730, Beijing 100080, China;

Institute of High Energy Physics, CAS, P.O. Box 918(4), Beijing 100039, China

The color suppressed modes $B^0 \rightarrow \bar{D}^{(*)0}\eta^{(\prime)}$ are analyzed in perturbative QCD approach. We find that the dominant contribution is from the non-factorizable diagrams. The branching ratios calculated in our approach for $B^0 \rightarrow \bar{D}^{(*)0}\eta$ agree with current experiments. By neglecting the gluonic contribution, we predict the branching ratios of $B^0 \rightarrow \bar{D}^{(*)0}\eta'$ are at the comparable size of $B^0 \rightarrow \bar{D}^{(*)0}\pi^0$, but smaller than that of $B^0 \rightarrow \bar{D}^{(*)0}\eta$.

PACS numbers: 13.25.Hw, 12.38.Bx

I. INTRODUCTION

The hadronic decays $B^0 \rightarrow \bar{D}^{(*)0}\eta^{(\prime)}$ are color suppressed modes, which belong to class II decays in the factorization approach (FA) [1]. The relevant effective weak Hamiltonian for these decays is given by

$$\mathcal{H}_{\text{eff}} = \frac{G_F}{\sqrt{2}} V_{cb} V_{ud}^* \left[C_1(\mu) O_1(\mu) + C_2(\mu) O_2(\mu) \right], \quad (1)$$

where the four-quark operators are

$$O_1 = (\bar{d}b)_{V-A} (\bar{c}u)_{V-A}, \quad O_2 = (\bar{c}b)_{V-A} (\bar{d}u)_{V-A}, \quad (2)$$

with the definition $(\bar{q}_1 q_2)_{V-A} \equiv \bar{q}_1 \gamma_\mu (1 - \gamma_5) q_2$. The Wilson coefficients $C_1 \sim -0.2$ and $C_2 \sim 1$ are calculated at m_b scale. The main contribution of these decays in FA is proportional to the Wilson coefficients $a_2 = C_1 + C_2/3$, which is a small number. That is the reason why class II decays usually have small branching ratios. Theoretical study of $B^0 \rightarrow \bar{D}^{(*)0}\eta^{(\prime)}$ decays gives a branching ratio of 10^{-5} [2]. However, recent experiments by Belle and BABAR show that the branching ratios of class II decays are not so small [3, 4]. The branching ratios of $B^0 \rightarrow \bar{D}^{(*)0}\eta$ are of 10^{-4} . Although the gluonic mechanism can enhance the $B^0 \rightarrow \bar{D}^0\eta'$ decay branching ratio to 10^{-4} [5], it may be difficult to explain the large branching ratio of $B^0 \rightarrow \bar{D}^{(*)0}\eta$. It means that the non-factorizable contributions in these decays are very important. This is confirmed in the recent theoretical study on charmed final state B meson decays in perturbative QCD approach [6].

The perturbative QCD approach (PQCD) for the exclusive hadronic B decays was developed some time ago [7, 8], and applied to the semi-leptonic [9] and non-leptonic decays [10, 11, 12] successfully. In this formalism, factorizable contributions, non-factorizable and annihilation contributions are all calculable. By including the k_T dependence of the wave functions and Sudakov form factor, this approach is free of endpoint singularity. Recent study shows that PQCD approach works well for charmless B decays [10, 11, 12], as well as for channels with one charmed meson in the final states [6, 13]. We will show the PQCD calculation of $B^0 \rightarrow \bar{D}^{(*)0}\eta^{(\prime)}$ decays in the next section, and discuss the numerical results in section III. The conclusion is in section IV.

II. $B^0 \rightarrow \bar{D}^{(*)0}\eta^{(\prime)}$ DECAY AMPLITUDES IN PQCD

In two-body hadronic B decays, the two outgoing mesons are energetic. Each of the valence quarks inside these mesons carries large momentum. Most of the energy comes from the heavy b quark decay in quark level. The light quark (d quark) inside B^0 meson, which is usually called spectator quark, carries small momentum at order of Λ_{QCD} . This quark also goes into final state meson in spectator diagrams. Therefore, we need an energetic gluon to connect this quark to the four quark operator involved in the b quark decay. Such that the spectator quark get energy from the four quark operator to form a fast moving light meson. The hard four quark dynamic together with the spectator quark becomes six-quark effective interaction. Since six-quark interaction is hard dynamics, it is perturbatively calculable. The non-perturbative dynamics in this process is described by the wave functions of mesons consisting of quark and anti-quark pair. The decay amplitude is then expressed as

$$\text{Amplitude} \sim \int d^4 k_1 d^4 k_2 d^4 k_3 \text{Tr} [C(t) \Phi_B(k_1) \Phi_{D^{(*)}}(k_2) \Phi_{\eta^{(\prime)}}(k_3) H(k_1, k_2, k_3, t) e^{-S(t)}]. \quad (3)$$




FIG. 1: Color-suppressed emission diagrams contributing to the $B^0 \rightarrow \bar{D}^{(*)0} \eta^{(')}$ decays.

Here $C(t)$ is the QCD corrected Wilson coefficient of the relevant four quark operator at scale t . Although next-to leading order results have been given [14], we will use leading order one here [11]. Φ_i are the meson wave functions, which include the non-perturbative contributions in these decays. The non-perturbative wave functions are not calculable in principal. But they are universal for all the hadronic decays. We will use the ones determined from other measured decay channels [6, 10, 11, 12]. The exponential $S(t)$ is the so-called Sudakov form factor, which includes the double logarithm resulting from the resummation of soft and collinear divergence. This form factor is also calculated to next-to leading order in the literature [15]. The Sudakov factor effectively suppresses the soft contributions in the process [6, 10, 11], thus it makes the perturbative calculation of hard part reliable.

Now the only left part of the decay amplitude is the hard part $H(t)$. Since it involves the four quark operator and the spectator quark connected by a hard gluon, it is channel dependent, but perturbatively calculable. There are altogether 8 kinds of diagrams in our $B^0 \rightarrow \bar{D}^{(*)0} \eta^{(')}$ decays, which are shown in Fig.1 for the spectator diagrams and Fig.2 for the annihilation type diagrams. Notice that in Fig.1, the $\eta^{(')}$ meson consists of $d\bar{d}$ content, while in Fig.2, it is a $u\bar{u}$ pair making $\eta^{(')}$ meson. Since $\eta^{(')}$ meson is isospin singlet ($u\bar{u} + d\bar{d}$), these two sets of diagrams give relatively positive contributions. On the other hand, in case of $B^0 \rightarrow \bar{D}^{(*)0} \pi^0$ decays [6], where π^0 is isospin triplet ($u\bar{u} - d\bar{d}$), these two sets of diagrams give destructive contributions there. Fortunately, as we will see later in the next section, the annihilation type diagrams are suppressed comparing to the spectator diagrams. Therefore, the branching ratios of these two kinds of decays are still comparable.

The structures of the meson wave functions are

$$B_{in}(P) : [\not{P} + m_B] \gamma_5 \phi_B(x), \quad (4)$$

$$D_{out}(P) : \gamma_5 [\not{P} + m_D] \phi_D(x), \quad (5)$$

$$D_{out}^*(P) : \not{\epsilon} [\not{P} + m_{D^*}] \phi_{D^*}(x), \quad (6)$$

$$\eta_{out}^{(')}(P) : \gamma_5 [\not{P} \phi_A(x) + m_0 \phi_P(x) + \zeta m_0 (\not{n}_- \not{n}_+ - 1) \phi_T(x)], \quad (7)$$

with $m_0 \equiv m_\pi^2/(m_u + m_d) = 1.4$ GeV, utilizing isospin symmetry. And the light-like vectors are defined as $n_+ = (1, 0, \mathbf{0}_T)$ and $n_- = (0, 1, \mathbf{0}_T)$. As shown in ref.[9], ϕ_B is identified as ϕ_+ and the contribution of another B meson wave function $\phi_B \propto \phi_+ - \phi_-$ is smaller in the PQCD calculations, therefore we neglect it. Applying for heavy quark symmetry, there is only one independent distribution amplitude $\phi_{D^{(*)}}$ in the heavy $D^{(*)}$ meson wave function [6, 13]. However, there are three distribution amplitudes for the light $\eta^{(')}$ meson wave functions [12], like the π meson wave function. The coefficients $\zeta = +1$ are for $\eta_{out}^{(')}$ with $\bar{u}$ ($\bar{d}$) carrying the momentum $x_3 P_3$, while $\zeta = -1$ for $\eta_{out}^{(')}$ with u (d) carrying the momentum $x_3 P_3$.

The gluonic mechanism of η' may make sizable contributions in the $B \rightarrow D^{(*)} \eta'$ decays, but they are usually model dependent [5]. Thus we will not consider it here. The η and η' mesons are mixtures of flavor SU(3) octet (η_8) and singlet (η_0) states in a two-mixing-angle formalism [16],

$$\begin{aligned} \eta &= \cos \theta_8 |\eta_8\rangle - \sin \theta_0 |\eta_0\rangle, \\ \eta' &= \sin \theta_8 |\eta_8\rangle + \cos \theta_0 |\eta_0\rangle. \end{aligned} \quad (8)$$

The definitions of the decay constants of η and η' are as follows:

$$\langle 0 | \bar{u} \gamma_\mu \gamma_5 u | \eta^{(')}(p) \rangle = i f_{\eta^{(')}}^u p_\mu, \quad \langle 0 | \bar{d} \gamma_\mu \gamma_5 d | \eta^{(')}(p) \rangle = i f_{\eta^{(')}}^d p_\mu. \quad (9)$$

The $\bar{s}s$ components of η and η' are not relevant in our decay channels. Therefore we did not show them. The decay constants in the two-angle mixing formalism are

$$f_\eta^u = f_\eta^d = \frac{f_8}{\sqrt{6}} \cos \theta_8 - \frac{f_0}{\sqrt{3}} \sin \theta_0, \quad (10)$$

$$f_{\eta'}^u = f_{\eta'}^d = \frac{f_8}{\sqrt{6}} \sin \theta_8 + \frac{f_0}{\sqrt{3}} \cos \theta_0. \quad (11)$$




FIG. 2: Annihilation Diagrams contributing to the $B^0 \rightarrow \bar{D}^{(*)0} \eta^{(')}$ decays.

The parameters are determined to be [16]

$$\begin{aligned} \theta_8 &= -22^\circ \sim -21^\circ, \quad f_8 = 1.28 f_\pi, \\ \theta_0 &= -9^\circ \sim -4^\circ, \quad f_0 = (1.20 - 1.25) f_\pi. \end{aligned} \quad (12)$$

The η and η' meson can also be expressed as a mixing in the quark flavor basis [17]

$$\begin{pmatrix} \eta \\ \eta' \end{pmatrix} = \begin{pmatrix} \cos \alpha & -\sin \alpha \\ \sin \alpha & \cos \alpha \end{pmatrix} \begin{pmatrix} (u\bar{u} + d\bar{d})/\sqrt{2} \\ s\bar{s} \end{pmatrix}, \quad (13)$$

where $\alpha = \pi + \theta - \arctan(1/\sqrt{2})$ describe the deviation from ideal mixing. The angle θ is one-angle mixing parameter. From the above eqn. (13), applying isospin symmetry, we have

$$f_\eta^u = f_\eta^d = f_\pi \cos \alpha / \sqrt{2}, \quad (14)$$

$$f_{\eta'}^u = f_{\eta'}^d = f_\pi \sin \alpha / \sqrt{2}. \quad (15)$$

The range of mixing parameters are determined to be $\theta = -17^\circ \sim -11^\circ$ [17].

The $B^0 \rightarrow \bar{D}^{(*)0} \eta^{(')}$ decay rate has the expression as,

$$\Gamma = \frac{1}{128\pi} G_F^2 |V_{cb}|^2 |V_{ud}|^2 m_B^3 |\mathcal{M}|^2. \quad (16)$$

Including the hard part and the meson wave functions, the $B^0 \rightarrow \bar{D}^{(*)0} \eta^{(')}$ decay amplitude is written as

$$\mathcal{M}(B^0 \rightarrow \bar{D}^{(*)0} \eta^{(')}) = f_{D^{(*)}} \xi_{\text{int}} + f_B \xi_{\text{exc}} + \mathcal{M}_{\text{int}} + \mathcal{M}_{\text{exc}}, \quad (17)$$

where $f_B = 190$ MeV, $f_D = f_{D^*} = 240$ MeV are the B and $D^{(*)}$ meson decay constants, respectively. The functions ξ_{int} , and ξ_{exc} denote the internal W -emission, and W -exchange contributions, which come from Figs. 1(a) and 1(b), Figs. 2(a) and 2(b), respectively. The functions $\mathcal{M}_{\text{int}}$, and $\mathcal{M}_{\text{exc}}$ represent the internal W -emission, and W -exchange contributions, which come from Figs. 1(c) and 1(d), Figs. 2(c) and 2(d), respectively. The expressions of the four functions are already shown in the appendix of ref.[6] for $B \rightarrow D\pi$ decays. One need only replace the pion wave function by the $\eta^{(')}$ wave function in those expressions.

In FA, only factorizable contribution of ξ_{int} (Fig.1(a)(b)) has been considered. Since ξ_{int} is proportional to the small Wilson coefficient $a_2 = C_1 + C_2/3$, the branching ratios predicted in FA is smaller than the experiments. Now in PQCD approach, all the topologies, including both factorizable and nonfactorizable ones, and also annihilation type ones have been taken into account. In fact the non-factorizable contribution M_{int} , which is proportional to the large Wilson coefficient $C_2/3$ is the dominant contribution in the $B^0 \rightarrow \bar{D}^{(*)0} \eta^{(')}$ decays. The reason is that the two non-factorizable diagrams in Fig.1(c) and (d) do not cancel each other like the B to two light meson decays, where the distribution amplitudes of wave function are symmetric [10, 11]. The large difference of $\bar{c}$ and u quark mass makes the contribution of M_{int} large. Very recently, the soft collinear effective theory also confirms that the non-factorizable M_{int} dominate over the contribution of $f_{D^{(*)}} \xi_{\text{int}}$ [18].

III. NUMERICAL RESULTS

As stated in the above section, we need various wave functions in our numerical calculations. Considering the previously calculations of other decay channels [6, 9, 10, 11, 12], the B meson wave function has been determined as

$$\phi_B(x, b) = N_B x^2 (1-x)^2 \exp \left[-\frac{1}{2} \left(\frac{x M_B}{\omega_B} \right)^2 - \frac{\omega_B^2 b^2}{2} \right], \quad (18)$$

TABLE I: PQCD predictions with one angle mixing formalism (I) and two angle mixing formalism (II) and experimental data (in units of 10^{-4}) of the $B^0 \rightarrow \bar{D}^{(*)0}\eta^{(\prime)}$ branching ratios.

Decay mode	PQCD (I)	PQCD (II)	Belle	BABAR	PDG
$B^0 \rightarrow \bar{D}^0\eta'$	$1.7 \sim 2.3$	$2.2 \sim 2.6$	-	-	< 9.4
$B^0 \rightarrow \bar{D}^0\eta$	$2.4 \sim 3.0$	$2.6 \sim 3.2$	$1.4^{+0.6}_{-0.5}$	2.41 ± 0.50	
$B^0 \rightarrow \bar{D}^0\pi^0$	2.3 ± 0.1		3.1 ± 0.6	2.89 ± 0.48	
$B^0 \rightarrow \bar{D}^{*0}\eta'$	$2.0 \sim 2.7$	$2.6 \sim 3.2$		-	< 14
$B^0 \rightarrow \bar{D}^{*0}\eta$	$2.8 \sim 3.5$	$3.1 \sim 3.8$	$2.0^{+1.0}_{-0.9}$	-	
$B^0 \rightarrow \bar{D}^{*0}\pi^0$	2.8 ± 0.1		2.7 ± 0.9	-	

where the shape parameter is chosen as $\omega_B = 0.4$ GeV. The normalization constant N_B is related to the decay constant f_B through

$$\int dx \phi_B(x, 0) = \frac{f_B}{2\sqrt{6}}. \quad (19)$$

The $D^{(*)}$ meson distribution amplitude is given by

$$\phi_{D^{(*)}}(x) = \frac{3}{\sqrt{6}} f_{D^{(*)}} x(1-x)[1 + C_{D^{(*)}}(1-2x)], \quad (20)$$

with the shape parameter $C_D = C_{D^*} = 0.8 \pm 0.2$ [6]. The range of $C_{D^{(*)}}$ was extracted from the $B \rightarrow D^{(*)}l\bar{\nu}$ decay spectrum at large recoil assuming $\omega_B = 0.4$ GeV for the B meson wave function [9]. We do not consider the variation of $\phi_{D^{(*)}}$ with the impact parameter b , since the current data are not yet sufficient to control this dependence. The light $\eta^{(\prime)}$ meson wave functions are chosen to be the same as the pion wave function according to isospin symmetry, since the relevant valence quarks here are mainly $u\bar{u}$ and $d\bar{d}$.

In the numerical analysis we adopt

$$\Lambda_{\overline{\text{MS}}}^{(f=4)} = 250 \text{ MeV}, \quad M_B = 5.2792 \text{ GeV}, \quad M_W = 80.41 \text{ GeV}.$$

Choosing $|V_{cb}| = 0.043$ and $|V_{ud}| = 0.974$, we obtain the PQCD predictions for the $B^0 \rightarrow \bar{D}^{(*)0}\eta^{(\prime)}$ branching ratios shown in Table I. For comparison, we also list the $B^0 \rightarrow \bar{D}^{(*)0}\pi^0$ decay branching ratios in this table. The theoretical uncertainty comes from the variation of the shape parameter for the $D^{(*)0}$ meson distribution amplitude, $0.6 < C_{D^{(*)}} < 1.0$ and the η and η' mixing parameter $\theta = -17^\circ \sim -11^\circ$ for the one-angle mixing formalism (I). The range of parameters of two-angle formalism (II) are shown in eq.(12). From numerical study, we notice that the branching ratios do not vary much upon the variation of $C_{D^{(*)}}$. This can be seen from the numbers of $B^0 \rightarrow \bar{D}^{(*)0}\pi^0$ in Table I, since it only depends on this parameter. Most of the uncertainty of $B^0 \rightarrow \bar{D}^{(*)0}\eta^{(\prime)}$ decays is from the mixing parameter θ or θ_8 and θ_0 . The branching ratios of $B^0 \rightarrow \bar{D}^{(*)0}\eta'$ increase while $B^0 \rightarrow \bar{D}^{(*)0}\eta$ decrease as the angle θ getting larger. Other input parameters, such as parameters of B meson wave function also affect the branching ratios, but they are mostly constrained by other well measured decay channels, like $B \rightarrow \pi\pi$ [11] and $B \rightarrow K\pi$ [10] decays, etc. The uncertainties of PQCD approach itself mainly come from the unknown higher twist contributions and higher order calculations of α_s corrections. No numerical estimation of higher twist contribution exist, although it is expected to be suppressed. The higher order QCD calculation in $B \rightarrow \phi K$ decay shows that next-to leading order α_s correction may not be small in certain channels [19].

The recently observed class-II decay $B^0 \rightarrow \bar{D}^{(*)0}\eta$ branching ratios are also listed in Table I [3, 4, 20]. It is easy to see that, our results agree with the experimental measurements within errors. Since we do not consider the extra gluon fusion contribution to $B^0 \rightarrow \bar{D}^{(*)0}\eta'$ decay, the not yet measured $B^0 \rightarrow \bar{D}^{(*)0}\eta'$ branching ratios are a little smaller than the $B^0 \rightarrow \bar{D}^{(*)0}\eta$ branching ratios. But they are still comparable with the $B^0 \rightarrow \bar{D}^{(*)0}\pi^0$ branching ratios. The reason for this comparable result is that we apply the assumption of exact isospin symmetry. We use the same wave function for $\eta^{(\prime)}$ and π meson where the only difference is the decay constant. The difference of dynamics is the constructive or destructive contribution from annihilation type diagrams. This destructive contribution makes the $B^0 \rightarrow \bar{D}^{(*)0}\pi^0$ branching ratios smaller than that of $B^0 \rightarrow \bar{D}^{(*)0}\eta$ decays. The numerical results also show that the dominant contribution comes from the non-factorizable contribution M_{int} . The factorizable contribution $f_D\xi_{int}$ and annihilation contribution M_{exc} are only 20-30% of M_{int} . Factorizable annihilation contribution $f_B\xi_{exc}$ is negligible.

IV. SUMMARY

In this work, we calculate the branching ratios of $B^0 \rightarrow \bar{D}^{(*)0}\eta^{(\prime)}$ decays in the perturbative QCD approach with k_T factorization, which is free of endpoint singularity. Belonging to class II decays in FA, these decays receive dominant contributions from the non-factorizable diagrams. Naive factorization breaks down in these color suppressed modes. The branching ratios calculated in our approach for $B^0 \rightarrow \bar{D}^{(*)0}\eta$ agree with current experiments. We predict the branching ratios of $B^0 \rightarrow \bar{D}^{(*)0}\eta'$ without gluonic contributions are at the comparable size of $B^0 \rightarrow \bar{D}^{(*)0}\pi^0$, but smaller than $B^0 \rightarrow \bar{D}^{(*)0}\eta$. They may be measured soon in the B factories.

Acknowledgments

We thank E. Kou, H.n. Li and A.I. Sanda for discussions. This work was supported by National Science Foundation of China under Grants No. 90103013 and 10135060.

-
- [1] M. Bauer, B. Stech, M. Wirbel, Z. Phys. C **29**, 637 (1985); *ibid.* **34**, 103 (1987); “NONLEPTONIC WEAK DECAYS OF B MESONS”, M. Neubert, B. second edition of “Heavy Flavours”, ed. by A.J. Buras and M. Lindner, World Scientific, Singapore, 1998.
 - [2] A. Deandrea and A.D. Polosa, Eur. Phys. J. C22, 677 (2002).
 - [3] Belle Colla., K. Abe *et al.*, Phys. Rev Lett. **88**, 052002 (2002).
 - [4] BABAR Colla., B. Aubert *et al.*, hep-ex/0207092.
 - [5] J.O. Eeg, A. Hiorth, A.D. Polosa, Phys. Rev. D65, 054030 (2002).
 - [6] H. Hayakawa, K. Hosokawa, T. Kurimoto, e-Print hep-ph/0212095; Y.-Y. Keum, T. Kurimoto, H. N. Li, C.-D. Lu, A.I. Sanda, e-Print Archive: hep-ph/0305335, Submitted to Phys. Rev. D.
 - [7] G.P. Lepage and S.J. Brodsky, Phys. Lett. B **87**, 359 (1979); G.P. Lepage and S.J. Brodsky, Phys. Rev. D **22**, 2157 (1980).
 - [8] J. Botts and G. Sterman, Nucl. Phys. **B225**, 62 (1989); H-n. Li and G. Sterman, Nucl. Phys. **B381**, 129 (1992).
 - [9] T. Kurimoto, H-n. Li, and A.I. Sanda, phys. Rev. D **65**, 014007, (2002); Phys. Rev. D67, 054028 (2003). C. D. Lü and M. Z. Yang, Eur. Phys. J. C28, 515 (2003).
 - [10] Y.Y. Keum, H-n. Li, and A.I. Sanda, Phys. Lett. B **504**, 6 (2001); Phys. Rev. D **63**, 054008 (2001); Y.Y. Keum and H-n. Li, Phys. Rev. **D63**, 074006 (2001).
 - [11] C. D. Lü, K. Ukai, and M. Z. Yang, Phys. Rev. D **63**, 074009 (2001).
 - [12] E. Kou and A. I. Sanda Phys. Lett. B525, 240 (2002).
 - [13] K. Ukai, p287, Preceedings for 4th International Conference on B Physics and CP Violation (BCP4), Japan, 2001; C-D Lü, Eur. Phys. J. C24, 121 (2002); hep-ph/0305061; C-D Lü and K. Ukai, Eur. Phys. J. C28, 305 (2003); Y. Li and C-D Lü, hep-ph/0304288; hep-ph/0305278.
 - [14] For a review, see G. Buchalla, A.J. Buras, M.E. Lautenbacher, Rev. Mod. Phys. **68**, 1125 (1996).
 - [15] H-n. Li and H.L. Yu, Phys. Rev. Lett. **74**, 4388 (1995); Phys. Lett. B **353**, 301 (1995); Phys. Rev. D **53**, 2480 (1996).
 - [16] H. Leutwyler, hep-ph/9709408, Nucl. Phys. Proc. Suppl. 64, 223 (1998); P. Herrera-Sikoldy, J.I. Latorre, P. Pascual and J. Taron, Phys. Lett. B419, 326 (1998); T. Feldmann and P. Kroll, Eur. Phys. J. C5 (1998) 327.
 - [17] T. Feldmann, P. Kroll, B. Stech, Phys. Rev. D58, 114006 (1998); E. Kou, Phys. Rev. D63, 054027 (2001).
 - [18] S. Mantry, D. Pirjol and I. Stewart, hep-ph/0306254.
 - [19] S. Mishima, A.I. Sanda, hep-ph/0305073.
 - [20] K. Hagiwara *et al.* (Particle Data Group), Phys. Rev. D **66**, 010001 (2002).